(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization International Bureau





(43) International Publication Date 30 October 2003 (30.10.2003)

PCT

(10) International Publication Number WO 03/089672 A1

(51) International Patent Classification7: C12Q 1/70, G01N 33/53, C12N 9/00, C07H 21/02, 21/04

(21) International Application Number: PCT/US03/12509

11 April 2003 (11.04.2003) (22) International Filing Date:

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data: 60/372,847

16 April 2002 (16.04.2002)

(71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): FLORES, Osvaldo, A. [CL/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). GROBLER, Jay [US/US]; 126 East Lincoln

Avenue, Rahway, NJ 07065-0907 (US). MURRAY, Edward, M. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). ZUCK, Paul, D. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(74) Common Representative: MERCK & CO., INC.; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(81) Designated States (national): CA, JP, US.

(84) Designated States (regional): European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR).

Published:

with international search report

before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: HEPATITIS C VIRUS ASSAY SYSTEMS

(57) Abstract: The present invention features assays employing a beta-lactamase reporter system, an HCV replicon enhanced cell, and/or a chimeric HCV replicon containing a 3' UTR based on the HCV-1a 3' UTR. These features can be employed alone or together, and are preferably combined together to measure HCV replicon activity and the affect of compounds on such activity